# Graded Brackets in Classical Field Theory: Conservative and Dissipative

XXXIII International Fall Workshop on Geometry and Physics

Rubén Izquierdo-López, joint work with M. de León and X. Rivas 9-13 September, 2025

**ICMAT-UNIR** 

### Introduction to the problem

Conserved quantities

Brackets in Classical Mechanics

### Geometric stage and algebraic structure of observables

Structure of the equations

Graded Poisson brackets

### Hamiltonians and extensions of brackets

### Into a theory of brackets in dissipative field theories


The Herglotz variational principle

Graded Jacobi brackets

### Conclusions and references

## Introduction to the problem


## Conserved quantities in field theories I



For 
$$\alpha \in \Omega^{n-1}(M)$$
:

$$\int_{X_{t_1}} \alpha = \int_{X_{t_2}} \alpha - \int_{\partial X \times [t_1, t_2]} \alpha \iff d\alpha = 0.$$

## Conserved quantities in field theories II



For 
$$\alpha \in \Omega^a(M)$$
:

$$\int_{X_{t_1}} \alpha = \int_{X_{t_2}} \alpha - \int_{\partial X \times [t_1,t_2]} \alpha \iff \mathrm{d}\alpha = 0.$$

### Proposed problem

Problem (that we would like to solve): Find all conserved quantities  $\sim$  Find forms that are closed on solutions.

**Problem (that we solve)**: Determine evolution of forms via some bracket:

$$\psi^*(d\alpha) = d\alpha + \{\alpha, \mathcal{H}\},\,$$

where  $\psi$  is a solution of the corresponding PDE and  $\alpha$  is some  $a\text{-}\mathrm{form}.$ 

### **Previous work:**

- Igor V. Kanatchikov. "Canonical Structure of Classical Field Theory in the Polymomentum Phase Space". In: Rep. Math. Phys. 41.1 (1998), pp. 49–90
- Miguel Á. Berbel and Marco Castrillón-López.
   "Poisson-Poincaré Reduction for Field Theories".
   In: J. Geom. Phys. 191 (2023), p. 104879
- François Gay-Balmaz, Juan C. Marrero, and Nicolás Martínez-Alba. "A New Canonical Affine Bracket Formulation of Hamiltonian Classical Field Theories of First Order". In: Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat. 118.3 (2024), p. 103

### Poisson bracket of functions

The Poisson bracket on the **phase space**  $T^*Q$  is defined as

$$\dot{f} = \frac{\partial f}{\partial t} + \{f, H\} = \frac{\partial f}{\partial t} + \frac{\partial f}{\partial q^i} \frac{\partial H}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial H}{\partial q^i},$$

where H is the Hamiltonian determining the dynamics.

Then, a time independent observable  $f \in C^{\infty}(M)$  is a conserved quantity if it is in **involution**  $\{f, H\} = 0$ .

On an arbitrary symplectic manifold  $(M, \omega)$ , where  $\omega \in \Omega^2(M)$ ,  $d\omega = 0$ , non-degenerate, is:

$$\forall f \in C^{\infty}(M), \exists ! X_f \in \mathfrak{X}(M), \iota_{X_f}\omega = \mathrm{d}f,$$

$$\{f,g\} := \omega(X_f,X_g) = X_g(f) = -X_f(g).$$

7

### **Poisson Theorem**

**Poisson Theorem**: If  $0 = \{f, H\} = \{g, H\}$ , we have  $\{\{f, g\}, H\} = 0$ .



Siméon Denis Poisson 1781-1840

### Jacobi theorem

### Jacobi Theorem:

$${f,{g,h}} + {h,{f,g}} + {g,{h,f}} = 0.$$



## Geometric stage and algebraic structure of observables

## General setup for field equations

Let  $\tau: M \longrightarrow X$  denote a fibered manifold,  $n = \dim X$ . Let:

- (i)  $\alpha_1, \ldots, \alpha_k \in \Omega^{n-1}(M)$  be semi-basic forms (representing observables).
- (ii)  $\beta_1, \ldots, \beta_k \in \Omega^n(M)$  be **semi basic forms** (representing evolution of observables).

**Semi-basic**: Locally, a semi-basic form is written as  $f dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_a}$ ,  $f \in C^{\infty}(M)$ .

We deal with partial differential equations with the following structure\*:

 $\psi^*(\mathrm{d}\alpha_i) = \beta_i \circ \psi$ , where  $\psi: X \to M$  is a subsection.

## **Examples (equations) I**

### Hamilton's equations:

- (a) Fiber bundle:  $T^*Q \times \mathbb{R} \to \mathbb{R}$ .
- (b) Semi-basic forms:  $q^i$ ,  $p_i$ , t.
- (c) Basic forms:  $\frac{\partial H}{\partial p_i} dt$ ,  $-\frac{\partial H}{\partial q^i} dt$ , dt.

And we obtain the equations:

$$\dot{q}^{i}dt = \frac{\partial H}{\partial p_{i}}dt$$
,  $\dot{p}_{i}dt = -\frac{\partial H}{\partial q^{i}}dt$ .

## **Examples (equations) II**

**Notation**: 
$$d^n x = dx^0 \wedge \cdots \wedge dx^{n-1}$$
,  $d^{n-1} x_\mu = (-1)^\mu dx^0 \wedge \cdots \wedge \widehat{dx^\mu} \wedge \cdots \wedge dx^{n-1}$ .

### Hamilton-De Donder-Weyl equations:

- (a) Fiber bundle: (covariant phase space)  $\bigwedge_{1}^{n} Y / \bigwedge_{1}^{n} Y \to X$ .
- (b) Semi-basic forms:  $y^i d^{n-1} x_\mu$ ,  $p_i^\mu d^{n-1} x_\mu$ ,  $\frac{1}{n} x^\mu d^{n-1} x_\mu$ .
- (c) Basic forms:  $\frac{\partial H}{\partial p_i^{\mu}} d^n x$ ,  $-\frac{\partial H}{\partial y^i} d^n x$ ,  $d^n x$ .

And we obtain the equations:

$$\frac{\partial y^{i}}{\partial x^{\mu}} d^{n}x = \frac{\partial H}{\partial p_{i}^{\mu}} d^{n}x, \quad \frac{\partial p_{i}^{\mu}}{\partial x^{\mu}} d^{n}x = -\frac{\partial H}{\partial y^{i}} d^{n}x.$$

## **Examples (equations) III**

### Yang–Mills equations:

- (a) Fiber bundle:  $\operatorname{Im} \operatorname{leg}_{\mathcal{L}} \to X$ .
- (b) Semi-basic forms:

$$A_{\mu}^{i}\mathrm{d}^{n-1}x_{\nu}-A_{\nu}^{i}\mathrm{d}^{n-1}x_{\mu},\;F_{i}^{\mu\nu}\mathrm{d}^{n-1}x_{\nu},\;\tfrac{1}{n}x^{\mu}\mathrm{d}^{n-1}x_{\mu}.$$

(c) Basic forms:

$$\left(-F^i_{\mu\nu}+f^i_{jk}A^j_\nu A^k_\mu\right)\mathrm{d}^nx,\,\left(-f^i_{jk}F^{\mu\nu}_iA^k_\mu\right)\mathrm{d}^nx,\,\mathrm{d}^nx.$$

And we obtain the equations:

$$\begin{split} &\left(\frac{\partial A^i_\mu}{\partial x^\nu} - \frac{\partial A^i_\nu}{\partial x^\mu}\right) \mathrm{d}^n x = \left(-F^i_{\mu\nu} + f^i_{jk} A^j_\nu A^k_\mu\right) \mathrm{d}^n x \,, \\ &\frac{\partial F^{\mu\nu}_i}{\partial x^\mu} \mathrm{d}^n x = -f^i_{jk} F^{\mu\nu} A^k_\mu \mathrm{d}^n x \,. \end{split}$$

### **Some comments**

- (i) Mostly any PDE fits into the description described earlier. However, we will require further structure in the sequel.
- (ii) So far we have given a local description, and what we will describe will work globally by requiring the previous picture to hold patch-wise on any fibered manifold  $\tau\colon M\to X$ . More particularly, we work with the subbundle

$$S^n := \langle d\alpha_1, \dots, d\alpha_k \rangle \subseteq \bigwedge^n T^*M,$$

and a map  $\beta \colon S^n \longrightarrow \bigwedge T^*X$ .

## Algebraic structure of observables

The observables  $\alpha_1 \dots, \alpha_k \in \Omega^{n-1}(M)$  allow us to define the space of **Hamiltonian forms**:

$$\Omega_H^{n-1}(M) := \{ \alpha \in \Omega^{n-1}(M) \colon d\alpha \in \langle d\alpha_1, \dots d\alpha_k \rangle \}.$$

**Assumption 1**: There is a bracket  $\{\cdot, \cdot\}$  on the space of Hamiltonian forms satisfying the following properties:

- (i) It is skew-symmetric:  $\{\alpha, \beta\} = -\{\beta, \alpha\}$ .
- $(\mathrm{ii})$  It satisfies the Jacobi identity up to an exact term:

$$\{\alpha,\{\beta,\gamma\}\}+\{\beta,\{\gamma,\alpha\}\}+\{\gamma,\{\alpha,\beta\}\}=\text{exact form}\,.$$

- (iii) It vanishes on closed forms:  $d\alpha = 0 \implies {\alpha, \beta} = 0$ .
- (iv) There is a correspondence  $\alpha \mapsto X_{\alpha}$  such that  $\{\alpha, \beta\} = \iota_{X_{\beta}} d\alpha$ .

### Parenthesis: How to obtain such a bracket?

Suppose we work with a **first order** variational problem on the sections of a configuration fiber bundle  $\pi\colon Y\to X$  given by a Lagrangian density

$$\mathcal{L}: J^1\pi \to \bigwedge^n T^*X, \quad \mathcal{L}(\phi) = L(x^\mu, \phi^i, \frac{\partial \phi^i}{\partial x^\mu}) d^n x.$$

The action is

$$\mathcal{J}[\phi] = \int_{X} \mathcal{L}(\phi).$$

We then find the multisymplectic form  $\Omega_{\mathcal{L}} \in \Omega^{n+1}(J^1\pi)$ , which is closed and allows to define the bracket:

$$\forall \alpha \in \Omega_H^{n-1}(J^1\pi), \exists X_\alpha \in \mathfrak{X}(M), \iota_{X_\alpha}\Omega_{\mathcal{L}} = d\alpha,$$

$$\{\alpha,\beta\} := \iota_{X_{\alpha} \wedge X_{\beta}} \Omega_{\mathcal{L}}.$$

## **Examples (algebraic structure)**

(i) **Hamilton equations**: Non vanishing brackets:

$$\{q^i,p_j\}=\delta^i_j.$$

(ii) Hamilton–De Donder–Weyl equations: Non vanishing brackets:

$$\{y^i {\rm d}^{n-1} x_\mu, p_j^\nu {\rm d}^{n-1} x_\nu\} = \delta_j^i {\rm d}^{n-1} x_\nu \,.$$

(iii) Yang-Mills equations: Non vanishing brackets:

$$\{A^i_\mu\mathrm{d}^{n-1}x_\nu-A^i_\nu\mathrm{d}^{n-1}x_\mu,F^{\alpha\beta}_j\mathrm{d}^{n-1}x_\beta\}=\delta^i_j\delta^{\alpha\beta}_{\mu\nu}\mathrm{d}^{n-1}x_\beta\,.$$

### Hamiltonian forms of arbitrary order I

### **Definition**

We say that a form  $\alpha \in \Omega^a(M)$  is **special Hamiltonian** if there is a semi-basic form  $\beta \in \Omega^{a+1}(M)$  such that  $\psi^*(\mathrm{d}\alpha) = \beta \circ \psi$ , for every solution of the equations. The space of special Hamiltonian *a*-forms is denoted by  $\widetilde{\Omega}_H^a(M)$ .

### Remark

If  $\alpha \in \Omega^{n-1}(M)$  is special Hamiltonian,  $\alpha \in \Omega^{n-1}_H(M)$ .

# Theorem (de León, I.L. 2025) $\alpha \in \Omega^a(M)$ is special Hamiltonian if and only if $\alpha \wedge \varepsilon \in \Omega^{n-1}_H(M)$ , for every closed and basic $\varepsilon \in \Omega^{n-1-a}(M)$ .

## Hamiltonian forms of arbitrary order II

### **Definition**

A form  $\alpha \in \Omega_H^a(M)$  is called **Hamiltonian** if  $d\alpha \in \iota_{\bigwedge^{n-(a+1)}TM}\langle d\alpha_1,\ldots,d\alpha_k\rangle$ . We denote by  $\Omega_H^a(M)$  the space of Hamiltonian *a*-forms.

### **Proposition**

$$\widetilde{\Omega}_H^a(M) \subseteq \Omega_H^a(M)$$


### Theorem (de León, I.L. 2025)

There is an unique induced graded Poisson bracket

$$\Omega_H^a(M)\otimes\Omega_H^b(M)\to\Omega_H^{a+b-(n-1)}(M)$$

that maintains the properties of the original bracket of (n-1)-forms. Furthermore, special Hamiltonian forms define a subalgebra.

## **Summary**




# Hamiltonians and extensions of brackets

### Domain of definition of current brackets

$$\begin{cases} \psi^*(d\alpha) = d\alpha + \{\alpha, \mathcal{H}\} \\ \deg\{\alpha, \mathcal{H}\} = \deg\alpha + \deg\mathcal{H} - (n-1) \end{cases} \implies \deg\mathcal{H} = n.$$

But:



### First extension of the brackets I

### Theorem (de León, I.L. 2025)

There exists an unique extension of  $\{\cdot,\cdot\}$ 

$$\Omega^{n-1}_H(M)\otimes\Omega^{\mathfrak{a}}_H(M)[1]\to\Omega^{\mathfrak{a}}_H(M)[1]$$

for arbitrary  $a \ge 0$  that satisfies the properties of  $\{\cdot,\cdot\}$ .


**Assumption 2**: There exists a form  $\mathcal{H} \in \Omega^n(M)[1]$ , the Hamiltonian, such that

$$\psi^*(d\alpha) = d\alpha + \{\alpha, \mathcal{H}\},\,$$

for every solution  $\psi$  and  $\alpha \in \Omega_H^{n-1}(M)$ .

### First extension of the brackets II

### Current domain of definition:



## **Examples (Hamiltonians)**

(i) Hamilton equations:

$$\mathcal{H} = H \mathrm{d}t - p_i \mathrm{d}q^i .$$

(ii) Hamilton–De Donder–Weyl equations:

$$\mathcal{H} = H \mathrm{d}^n x - p_i^{\mu} \mathrm{d} y^i \wedge \mathrm{d}^{n-1} x_{\mu} \,.$$


(iii) Yang-Mills equations

$$\mathcal{H} = \left( -\frac{1}{4} F_i^{\mu\nu} F_{\mu\nu}^i + \frac{1}{2} f_{jk}^i F_i^{\mu\nu} A_{\mu}^j A_{\nu}^k \right) d^n x - F_i^{\mu\nu} dA_{\mu}^i \wedge d^{n-1} x_{\nu} .$$

Hamiltonian = Poincaré-Cartan form

### Final extension of the bracket I

Question: Can we interpret  $\psi^*(d\alpha) = d\alpha + \{\alpha, \mathcal{H}\}$ , for arbitrary  $\alpha \in \Omega^a_H(M)$ ?



### Final extensions of the brackets II

**Problem**: There is no unique extension of the bracket to order a < n - 1.

Nevertheless.

### Theorem (de León, I.L. 2025)

There is a bijective correspondence between the possible extensions of the bracket and affine maps

 $\gamma: \{\textit{Hamiltonians}\} \rightarrow \{\textit{Ehresmann connections on } \tau: \textit{M} \rightarrow \textit{X}\}$ 

such that  $\gamma(\mathcal{H})$  solves the Hamilton–De Donder–Weyl equations of  $\mathcal{H}$ , for every  $\mathcal{H}$ .

### Final extension of the bracket III

### Corollary

Let  $\gamma$  be such a map,  $\mathcal H$  be a Hamiltonian, and  $\psi$  be an integral section of  $\gamma(\mathcal H)$ . Then

$$\psi^*(\mathrm{d}\alpha) = \mathrm{d}\alpha + \{\alpha, \mathcal{H}\}_{\gamma},$$

for every  $\alpha \in \Omega_H^a(M)$ .

### **Corollary**

Let  $\alpha$  be a special Hamiltonian form. Then, the bracket  $\{\alpha, \mathcal{H}\}_{\gamma}$  is independent of extension  $\{\cdot, \cdot\}_{\gamma}$ , for every Hamiltonian  $\mathcal{H}$ .

### Theorem (Poisson theorem)

Special Hamiltonian forms that are closed on solutions close a subalgebra under the graded Poisson bracket.

### **Technical remarks**

The construction was based on a generalization of the  $\sharp$  mapping associated to a graded Poisson bracket. In particular, we generalized the techniques employed in

- Peter W. Michor. "A Generalization of Hamiltonian Mechanics". In: J. Geom. Phys. 2.2 (1985), pp. 67–82
- Janusz Grabowski. "Z-Graded Extensions of Poisson Brackets". In: Rev. Math. Phys. 09.01 (1997), pp. 1–27

to extend the brackets.

Into a theory of brackets in

dissipative field theories

## The Herglotz variational principle for fields I

We have presented the previous questions in the usual framework of variational problems. However, there is a growing interest in more general variational problems, such as those involving Lagrangians dependent on the action.

- Matheus J. Lazo et al. "An Action Principle for Action-Dependent Lagrangians: Toward an Action Principle to Non-Conservative Systems". In: J. Math. Phys. 59.3 (2018). 10.1063/1.5019936, p. 032902
- Jordi Gaset et al. "The Herglotz Variational Principle for Dissipative Field Theories". In: Geom. Mech. 01.02 (2024). 10.1142/S2972458924500060, pp. 153–178

## The Herglotz variational principle for fields II

Let  $\pi\colon Y\to X$  be a fiber bundle, and let  $\mathcal{L}\colon J^1\pi\times_X \bigwedge^{n-1} T^*X\to \bigwedge^n T^*X$  be a Lagrangian dependent on the action. Locally,

$$\mathcal{L} = L(x^{\mu}, \phi^{i}, \frac{\partial \phi^{i}}{\partial x^{\mu}}, s^{\mu}) d^{n}x.$$

We look for fields  $\phi^i(x^\mu)$  minimizing the following problem:

Look for 
$$\zeta^{\mu}$$
 satisfying  $\frac{\partial \zeta^{\mu}}{\partial x^{\mu}} = L\left(x^{\mu}\phi^{i}, \frac{\partial \phi^{i}}{\partial x^{\mu}}, \zeta^{\mu}\right)$ , Minimize  $\int_{X} L \mathrm{d}^{n}x$ .

## The Herglotz equations of motion and multicontact geometry

The equations obtained with the previous variational principle are the **Herglotz equations of motion**:

$$\begin{split} \frac{\partial L}{\partial \phi^i} - \frac{\mathrm{d}}{\mathrm{d} x^\mu} \left( \frac{\partial L}{\partial (\partial_\mu \phi^i)} \right) + \frac{\partial L}{\partial s^\mu} \frac{\partial L}{\partial (\partial_\mu \phi^i)} = 0 \,, \\ \frac{\partial s^\mu}{\partial x^\mu} = L \,. \end{split}$$

Intrinsically, these are described by the multicontact form  $\Theta_{\mathcal{L}} \in \Omega^n(J^1\pi)$ .

$$\Theta_{\mathcal{L}} = \left( L - \partial_{\mu} \phi^{i} \frac{\partial L}{\partial (\partial_{\mu} \phi^{i})} \right) d^{n} x + \frac{\partial L}{\partial (\partial_{\mu} \phi^{i})} d\phi^{i} \wedge d^{n-1} x_{\mu} + ds^{\mu} \wedge d^{n-1} x_{\mu}.$$

### Graded Jacobi brackets I

Let  $M := J^1 \pi \times_X \bigwedge^{n-1} T^* X$ . Borrowing from contact geometry:

We say that a multivector field  $X \in \mathfrak{X}^p(M)$  (sum of decomposable  $X_1 \wedge \cdots \wedge X_p$ ), is a **infinitesimal conformal** transformation of  $\Theta_{\mathcal{L}}$  if

$$\mathcal{L}_U \Theta_{\mathcal{L}} = d\iota_U \Theta_{\mathcal{L}} + (-1)^{p-1} \iota_U d\Theta_{\mathcal{L}} = \iota_V \Theta_{\mathcal{L}},$$

for certain  $V \in \mathfrak{X}^{p-1}(M)$  called the **conformal factor**.

#### Graded Jacobi brackets II

#### **Definition**

A form  $\alpha \in \Omega^a(M)$  is called **conformal Hamiltonian** if there is a conformal multivector field U such that  $\alpha = \iota_U \Theta$ . The space of conformal Hamiltonian forms is denoted by  $\Omega^a_H(M)$ .

## Theorem (de Léon, I.L., Rivas )

Let  $U_1$  and  $U_2$  be conformal infinitesimal transformations of  $\Theta_{\mathcal{L}}$ . Then,  $\iota_{[U_1,U_2]}\Theta_{\mathcal{L}}$ , where  $[U_1,U_2]$  is the Schouten–Nijenhuis bracket, only depends on the values of  $\iota_{U_1}\Theta_{\mathcal{L}}$  and  $\iota_{U_2}\Theta_{\mathcal{L}}$ .

#### Definition

Then,  $\{\alpha_1, \alpha_2\} := -\iota_{[U_1, U_2]} \Theta_{\mathcal{L}}$  is called the **graded Jacobi bracket** of  $\alpha_i := \iota_{U_i} \Theta_{\mathcal{L}}$ .

#### Graded Jacobi brackets III

Defining  $\deg_H \alpha := (n-1) - \deg \alpha$ , the bracket has the following properties:

**Theorem (de León, I.L., Rivas)**The graded Jacobi bracket defines an operation

$$\Omega_H^a(M)\otimes\Omega_H^b(M)\xrightarrow{\{\cdot,\cdot\}}\Omega_H^{a+b-(n-1)}(M)$$

satisfying the following properties:

- (i) It is graded-skew-symmetric  $\{\alpha, \beta\} = -(-1)^{\deg_H \alpha \deg_H \beta} \{\beta, \alpha\};$
- (ii) It satisfies the graded Jacobi identity:

$$(-1)^{\deg_H \alpha \deg_H \gamma} \{\alpha, \{\beta, \gamma\}\} + cycl. = 0.$$

# **Conclusions and references**

# Final remarks and remaining questions

- (i) The previous theoretical results seem to indicate that the subalgbra of special Hamiltonian forms is of high relevance to a particular field theory. We would like to compute these subalgebras for several almost regular Lagrangians to further study these classical field theories.
- (ii) (In progress) We would also like to investigate the relation between these extensions and the instantaneous split formalism.
- (iii) It is also interesting to investigate the implications of this algebraic structure in the study of momentum maps and reduction, employing the graded brackets.
- (iv) Is it possible to extend all of the results in the conservative scenario to the dissipative scenario?

#### Main references

- Manuel de León and Rubén Izquierdo-López. "Graded Poisson and Graded Dirac Structures". In: J. Math. Phys. 66.2 (2025). 10.1063/5.0243128, p. 022901
- Manuel de León and Rubén Izquierdo-López. A
   Description of Classical Field Equations Using Extensions of Graded Poisson Brackets.
   10.48550/arXiv.2507.04743. 2025
- Manuel de León, Rubén Izquierdo-López, and Xavier Rivas. *Brackets in Multicontact Geometry* and Multisymplectization. 10.48550/arXiv.2505.13224. 2025

Thank you for your attention!


# The graded Dirac structure of an almost regular Lagrangian

#### **Graded Dirac structures I**

#### **Definition**

A (fibered) **graded Dirac structure** on a fibered manifold  $\tau \colon M \to X$  is a tuple  $(S^a, \sharp_a)$ , where


- (i)  $S^a$ , a = 1, ..., n is a subbundle of  $\bigwedge^a T^*M$ , and is composed of (a-1)-horizontal forms.
- (ii)  $\sharp_a$  are vector bundle mappings



#### **Graded Dirac structures II**

These maps  $\sharp_a$  are required to satisfy the following

- (a) They are skew-symmetric:  $\iota_{\sharp_{a}(\alpha)}\beta = (-1)^{(n+1-\deg\alpha)(n+1-\deg\beta)}\iota_{\sharp_{\beta}}\alpha$
- (b) They are **integrable** (involutive with respect to a generalization of the Courant bracket).



# Idea relating to brackets

#### Graded Dirac structures ~ infinitesimal version of brackets

- (i)  $S^a$  is characterized as the subbundle such that  $\alpha \in \Omega_H^{a-1}(M)$  if and only if  $d\alpha \in S^a$ .
- (ii)  $\sharp^a$  is characterized as  $\{\alpha, \beta\} = (-1)^{\deg_H \beta} \iota_{\sharp_a(\mathrm{d}\alpha)} \mathrm{d}\beta$ .
- (iii) Involutivity is characterized by the Jacobi identity.

In fact,

#### **Theorem**

 $\{\mathit{Graded\ Poisson\ brackets}\}^\star\cong\{\mathit{Graded\ Dirac\ structures}\}$ 

# Basic constructions related to graded Dirac structures

## Theorem (de León, I.L. 2025)

- (i) **Pullbacks**:  $f: M_1 \rightarrow M_2$ ,  $M_2$  is graded Dirac, there is a graded Dirac structure on  $M_1$ .
- (ii) **Pushforwards**:  $\pi: M_1 \to M_2$  submersion,  $M_1$  graded Dirac, then\* there is a graded Dirac on  $M_2$ .

 $\star = Vertical vector field define symmetries$ 

# Almost regular Lagrangians I

Let  $\mathcal{L} = L(x^{\mu}\phi^{i}\partial_{\mu}\phi^{i}) d^{n}x$ . We define the **Legendre transformation** as the fibered derivative:


$$\operatorname{Leg}_{\mathcal{L}} \colon J^1 \pi \to \bigwedge^n T^* Y$$
.

There is a canonical **multisymplectic structure** on  $\bigwedge^n T^*Y$ , which is a closed, non-degenerate (n+1)-form  $\Omega$ .

We recover the multisymplectic form as  $\Omega_{\mathcal{L}} = \mathsf{Leg}_{\mathcal{L}}^* \Omega$ .

# Almost regular Lagrangians II

Generally, working on  $(J^1\pi, \Omega_{\mathcal{L}})$  is **hard**, and on  $\bigwedge^n T^*Y$  are too many variables. What is easier to work is the submanifold Im Leg<sub> $\mathcal{L}$ </sub> or, rather,the image under:



which inherits a natural graded Dirac structure.